

Physicochemical properties and quality evaluation of white bread with “Baromi 2” rice flour as a wheat flour substitute

Ga Hyeon Kim^{1,2,*}, Sung Huo Kim^{1,2,*}, Yeon-Ji Jo^{2,3} and Sung Hoon Park^{1,2,*}

¹Department of Food & Nutrition, College of Life Sciences, Gangneung-Wonju National University, Gangneung 25457, Korea

²Haeram Institute of Bakery Science, Gangneung-Wonju National University, Gangneung 25457, Korea

³Department of Marine Bio Food Science, Gangneung-Wonju National University, Gangneung 25457, Korea

Abstract

This study aimed to broaden the utilization of rice flour by incorporating it into white bread. The rice flour used was derived from the “Baromi 2 (B2)” soft rice variety developed by the Rural Development Administration. Wheat flour in white bread was substituted with B2 at levels of 20%, 40%, 60%, 80%, and 100%. The physicochemical properties of the resulting products were evaluated. As B2 substitution levels increased, fermentation expansion and product volume decreased. However, no significant differences from the control group were observed substitution levels up to 20%. Increasing B2 levels also reduced the crust's Browning Index (BI), whereas products with appropriate B2 substitution exhibited reduced staling rates. These findings suggest that B2 has the potential to be used as a wheat flour substitute at levels up to 20% in baked goods. This study also reaffirms B2's suitability as an alternative to wheat flour, suggesting that incorporating various baking enzymes and gluten substitutes will improve the quality of rice-based bread products in future applications.

Keywords: Baromi2, Floury rice, Lower gluten, Healthy bread

Introduction

Rice (*Oryza Sativa* L.) ranks among the most commonly eaten grains, used in various forms, including noodles and products like rice cakes, alcoholic drinks, sweets, and baked items (Chu et al., 2023). As eating habits become more varied and convenience grows, rice consumption continues to decline. Additionally, the increase in single-person and dual-income households has resulted in higher consumption of home meal replacements (KOSTAT, 2022). Transitioning to Westernized diets has led South Korea to maintain a structural surplus since 2000, as annual rice production and imports have repeatedly surpassed domestic consumption.

At the same time, individual rice consumption in South Korea

has declined by 17.3%, dropping from 69.8 kg in 2012 to 57.5 kg in recent years (KDCA, 2022). During the 1990s, per capita rice consumption dropped annually by an average of 2.3%, which rose to 2.6% in the 2000s (Kim, 2017). To encourage rice consumption, it is essential to expand its application beyond just a staple food to include its use as an ingredient in processed foods (Choi & Lee, 2007).

The rise in single-person households swiftly refocused domestic rice consumption from traditional table rice to processed alternatives (Ha et al, 2022). The rice processing industry's market has steadily increased from 4.1 trillion won in 2019 to 7.3 trillion won (KOSTAT, 2022), heightening interest in using surplus rice and rice-based processed foods. Ongoing research centers on bakery products that

Received: Nov 06, 2024 / Revised: Nov 19, 2024 / Accepted: Nov 19, 2024

Corresponding author: Sung Hoon Park, Department of Food & Nutrition, College of Life Sciences, Gangneung-Wonju National University, Gangneung 25457, Korea

E-mail: sungpark@gwnu.ac.kr

*These authors contributed equally to this work.

Copyright © 2024 Korean Society for Food Engineering.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

use these materials (Qin et al., 2021; Fujii et al., 2023).

Bread mainly relies on wheat flour due to the essential gluten proteins it contains (Aljada et al., 2021). Gluten is an alcohol-soluble protein in various grains, including wheat, rye, and barley (Caio et al., 2019), consisting of gliadin and glutenin. Gluten may cause several conditions, such as gluten sensitivity, celiac disease, gastrointestinal issues, and dermatitis herpetiformis, the latter presenting with skin symptoms associated with digestive problems (Shewry, 2019; Šmídová & Rysová, 2022). As a result, gluten-reduced diets are becoming increasingly popular for their purported health benefits (Cannas et al., 2020).

Recent research has investigated the feasibility of using soybeans, corn, and sorghum as alternatives to wheat in bread products (Sun et al., 2019; Woomer et al., 2019; Mariscal-Moreno et al., 2021). Nonetheless, research on using rice as a wheat flour replacement is still scarce (Kim, 2017). Unlike wheat, rice lacks gliadin, preventing the development of a gluten network needed for product cohesion (Honda et al., 2021). Additionally, rice flour quality declines during storage, potentially leading to lipid spoilage and rancidity (Zhao et al., 2021).

"Baromi 2 (B2)" is a new rice variety created by the Rural Development Administration. It results from crossing "Suwon 524," used for milling, with "Jopyeong," a versatile variety adapted to upland conditions bred for resistance (Ha et al., 2022). Rice is typically subjected to wet milling for powdered form due to its greater grain hardness than wheat, primarily processed through wet milling following starch gelatinization (Chu et al., 2023). B2 is a dry-milled rice flour that offers cost savings, shorter processing times, and environmental benefits over wet-milled rice flour (Park et al., 2023). In the food sector, powdered rice flour can be used in nonfermented bread, fish cakes with low wheat content, and sausages. Additionally, it can be combined with wheat flour to create fermented bread and other processed foods (Ministry of Agriculture, 2022).

As a result, this research aims to improve the use of B2 in bakery products to encourage rice flour consumption. It evaluates B2's potential as a wheat flour substitute in white bread while examining its physicochemical properties. Furthermore, the study aims to identify the ideal blending ratio of rice flour as a wheat flour replacement and offers essential data for rice flour processing studies.

Materials and Method

Materials

Rice flour was made from B2 supplied by the Rural Development Administration.

The patent application is named "Garumi2." The hardness value of "Garumi2" is 2.9 ± 0.09 kg. The physicochemical properties of the rice flour produced with the test mill "Buhler MLU-202" include average particle size (μm) 81.8 ± 1.31 , damaged starch (%) 5.6 ± 0.11 , whiteness (CIE value) 92.3 ± 0.22 , ash content (%) 0.59 ± 0.02 , protein content (%) 6.6 ± 0.16 , and amylose content (%) 18.0 ± 0.59 (Jung et al., 2020).

Preparation of white pan bread and dough

White pan bread samples were prepared using the AACC standard straight dough method with slight modifications (AACC, 2000). The ingredients were as follows: 100 g of hard flour (Daehan Flour Mills Co., Ltd, Incheon, Korea), 4 g of refined sugar (CJ. Inc., Seoul, Korea), 4 g of whole dry milk (Seoul Dairy Cooperative, Seoul, Korea), 6 g of shortening (Samyangsa Co. Inc., Incheon, Korea), 2 g of sea salt, and 0.7 g of yeast (Saf-instant gold, Lesaffre Pte Ltd, France) for the control (RC) sample. For the tests, 20% (R1), 40% (R2), 60% (R3), 80% (R4) and 100% (R5) concentrations of B2 rice flour were added instead of wheat flour (Table 1). We mixed the ingredients for 3 min at low speed and then for 7 min at high speed (SK-20 Mixer, SK Mixer Co., Ltd. Saitama, Japan). The mixed dough was allowed to ferment for 1 h at 27°C with 75%-80% humidity. After sheeting, molding, and proofing for 60 min at 38°C with 85%-95% humidity, the dough was baked at 170°C (upper heat) and 210°C (lower heat) for 35 min (Fig. 1). The finished products were cooled until their internal temperature reached 30°C before conducting the experiments.

Analysis of dough pH and fermentation gas retention capability

The dough's pH was measured using a pH meter after mixing 5 g of dough and 45 mL of distilled water (S20 SevenEasy, Mettler Toledo, Leicester, UK). The dough's fermentation expansion was assessed by placing 10 g of bread dough into a 50 mL conical tube, smoothing the surface evenly, and subjecting it to fermentation for

Table 1. Baker’s percentages of white pan bread ingredients

	RC	R1	R2	R3	R4	R5
Flour	100	80	60	40	20	
Baromi 2		20	40	60	80	100
Refined sugar	4	4	4	4	4	4
Whole dry milk	4	4	4	4	4	4
Salt	2	2	2	2	2	2
Instant dry yeast	0.7	0.7	0.7	0.7	0.7	0.7
Shortening	8	8	8	8	8	8
Water	68	68	68	68	68	68

The following table represents the ingredients used in white pan bread production expressed in baker’s percentages: RC-contains 0% “Baromi2 (B2)”; R1-contains 20% B2; R2-contains 40% B2; R3-contains 60% B2; R4-contains 80% B2; R5-contains 100% B2.




Fig. 1. Schematic diagram of the white pan bread-making process.

120 min at 20 min intervals (temperature 27°C, humidity 85%) when measurements were taken. All samples started fermentation at the same volume (9 mL); the highest point of dough expansion was measured every 20 min.

Analysis of baking loss rate, specific volume, and height of the product

The specific volume of the bread was measured 24 h after product molding using the VOLSCAN VSP300 (Stable Micro Systems Ltd, GU, UK) under the condition of 4 mm/sec. The baking loss rate was calculated using the following formula: the

bread dough’s weight and the bread’s weight. All samples were measured five times.

$$\text{Baking Loss Rate (\%)} = \frac{\text{Before baking (g)} - \text{After baking (g)}}{\text{Before baking (g)}} \times 100$$

Moisture content of the products

The finished bread’s moisture content was assessed using the Infrared Moisture Analyzer FD-720 (Kett, Tokyo, Japan). A 3.000±0.005 g test sample was placed in an 8 cm diameter aluminum pan. The drying process temperature was maintained at 105°C. Measurements lasted approximately 8 minutes, and results were expressed as a percentage. Each variant was tested in triplicate.

Color analysis of the products

The color measurement of the crust and crumb of white pan bread was conducted using a Chromameter CR-400 (Konica Minolta, Tokyo, Japan) with the CIE L*, a*, b* system. Calibration was performed with a standard white plate, and the measurement diameter was set at 0.8 cm. Each sample had five readings taken. The L* coordinate indicates surface brightness on a scale from 0 to 100, while a* measures color saturation, with positive values representing red and negative values representing green. Positive values denote yellowness for the b* parameter, and negative values indicate blueness. The crust’s Browning Index (BI) was calculated using the L*, a*, and b* parameters (Kupiec et al., 2021).

$$BI = \frac{[100 \times (x - 0.31)]}{0.172}$$

$$X = \frac{(a^* + 0.75 \times L^*)}{(5.645 \times L^* + a^* - 3.012 \times b^*)}$$

Texture profile analysis (TPA) of the products

The TPA for white pan bread and sweet bread was conducted using the CTX Texture Analyzer (AMETEK Brookfield, MA, USA) with these settings: table speed at 120 mm/min, a maximum load of 2 kg, a compression rate of 50%, and performed under 2-bite conditions. The samples analyzed were 20 mm thick slices taken from the center of the bread. Four parameters were assessed: hardness (N), springiness (mm), cohesiveness, and chewiness (J). Measurements were carried out at 24-, 48-, and 96-hr postmanufacturing.

Statistical analysis


The results were analyzed using the Statistical Package for the Social Sciences. ANOVA was applied, followed by the Tukey test for post hoc comparisons. The statistical significance level was $p < 0.05$.

Results and Discussion

Physicochemical quality characteristics of white pan bread and dough


The external structure and attributes of bread containing B2 are displayed in Table 2, Fig. 2, and Fig. 3. Fermentation expansion measurements for the B2-enriched bread products are depicted in Fig. 3. Over a 120-minute fermentation period, the dough volume increased by 23.83 mL for the control group (RC), while R1 showed

an increase of 25.33 mL, R2 saw an 18.00 mL rise, R3 increased by 10.5 mL, R4 by 7.25 mL, and R5 by 3.5 mL. The decline in fermentation expansion at higher B2 substitution levels is linked to the reduced gluten content, which is crucial for dough rising.



White pan bread crust (see image at the top – A) and its cross-section (see image at the bottom – B): RC-contains 0% B2, R1 contains 20% B2, R2 contains 40% B2, R3 contains 60% B2, R4 contains 80% B2, and R5 contains 100% B2.

Fig. 2. Cross-sectional analysis of white pan bread with “Baromi2”.



RC-contains 0% B2; R1-contains 20% B2; R2-contains 40% B2; R3-contains 60% B2; R4-contains 80% B2; R5-contains 100% B2. Each experiment was conducted in triplicate, and the results are presented as the mean \pm SD.

Fig. 3. Fermentation capacity of white pan bread with “Baromi2”.

Table 2. Physicochemical quality characteristics of white pan bread containing “Baromi2”.

	RC	R1	R2	R3	R4	R5
Volume (mL)	1,640.25 \pm 9.70 ^a	1,504.85 \pm 12.12 ^b	1,203.32 \pm 15.06 ^c	1,052.34 \pm 10.16 ^d	452.94 \pm 14.27 ^e	391.63 \pm 8.41 ^f
Weight (g)	321.08 \pm 1.68 ^a	321.38 \pm 1.92 ^a	324.12 \pm 1.00 ^a	322.69 \pm 1.56 ^{ab}	327.06 \pm 1.22 ^b	326.88 \pm 1.90 ^b
Height (mm)	117.46 \pm 1.48 ^{ab}	108.18 \pm 1.56 ^a	91.16 \pm 8.40 ^b	60.48 \pm 7.21 ^c	35.44 \pm 3.02 ^d	33.01 \pm 2.45 ^e
Specific volume (mL/g)	3.98 \pm 0.09 ^a	3.81 \pm 0.09 ^a	3.23 \pm 0.10 ^b	2.21 \pm 0.09 ^c	1.37 \pm 0.11 ^d	0.98 \pm 0.01 ^e
Baking loss rate (%)	15.51 \pm 0.44 ^a	15.43 \pm 0.50 ^a	14.71 \pm 0.26 ^b	15.08 \pm 0.41 ^b	13.93 \pm 0.32 ^c	13.98 \pm 0.50 ^c
Moisture content (%)	33.60 \pm 0.25 ^{ab}	32.59 \pm 0.51 ^a	34.37 \pm 0.40 ^b	33.03 \pm 0.78 ^{ab}	34.29 \pm 0.31 ^b	32.57 \pm 0.15 ^a

White pan bread's volume, weight, height, and baking loss rate are as follows: RC-contains 0% “Baromi2(B2),” R1 contains 20% B2, and R2 contains 40%.

^{a-f}Above each column signify a statistical difference at $\alpha=0.05$.

^{NS}Indicates non-significant results. Different letters.

Nonetheless, the data indicates that fermentation expansion can be enhanced at an optimal substitution level. Notably, substituting 20% B2 in the dough resulted in fermentation expansion exceeding that of the control group without negatively impacting the dough's physical properties (Jun et al., 2019).

Table 2 indicates that increasing the B2 content decreased the bread's volume and height, accompanied by a rise in weight. The moisture content analysis showed no substantial difference between RC and R1, although a trend was observed where moisture content tended to rise with a higher proportion of rice flour.

This is probably caused by the dough's diminished ability to retain gas, attributed to lower gluten development, resulting in lesser oven spring and heightened moisture retention (Kraithong et al., 2018). Consequently, the baking loss rate lessened with increased B2 content. The specific volume is a critical measure of bread quality (Monteiro et al., 2021). This study found that the volume of bread with B2 diminished as the substitution level of B2 rose. Additionally, the baking loss rates analysis indicated no significant difference between RC (15.42%) and R1 (14.47%), while R2 showed a lower value (14.01%).

Color analysis of white pan bread

The analysis focused on the bread's color and varying B2 levels to determine how increased B2 affects it (see Fig. 2, Table 3). Measurements of the crumb color indicated that as the B2 content rose, the L* value of the crumb also increased, with the * value remaining stable across all samples. The Tb* value showed a positive trend alongside the rising B2 content. In summary, while

the L* value of the crumb decreased, the b* value increased, suggesting that higher B2 content contributes to a darker crumb color. Conversely, as the B2 content rose, the L* value of the crust improved, while both the a* and b* values declined. The BI, derived from the crust's L*, a*, and b* values, demonstrated that the crust color lightened with increased B2 substitution levels. These findings indicate that the rate of B2 substitution for wheat flour notably influences the color of the final product.

Rice flour contains lower protein levels than wheat flour, which is naturally rich in carotenoids, contributing to its darker color (Choi, 2010; Baek & Koh, 2014). Changes in crust color can be linked to the higher B2 content. These results are consistent with earlier studies on rice flour-based bread, which indicated that products with rice flour had greater L* values than the control group. Furthermore, from sample R3 onward, the bread crumb displayed larger pores and inadequate fermentation, rendering these samples inappropriate for further analysis.

TPA of white pan bread

Texture parameters like hardness, cohesiveness, springiness, and chewiness are essential quality indicators for baked goods consumers (Roman et al., 2020). Among these factors, a rise in hardness is the most noticeable and frequently observed parameter linked to staling (Kim et al., 2019). The texture of the bread with B2 was evaluated 24 hr after baking. Hardness was measured after 4 days of storage, precisely at 24-, 48-, and 96-hr after baking (Table 4). At the 24-hr mark, the texture assessment revealed no significant differences in cohesiveness, springiness, or chewiness among the samples. Regard-

Table 3. Color analysis of white pan bread contained "Baromi2"

		RC	R1	R2	R3	R4	R5
Crumb	L*	77.51±1.45 ^a	79.52±0.99 ^b	77.59±1.39 ^a	73.90±1.19 ^c	75.07±0.87 ^{cd}	75.43±0.31 ^d
	a*	2.48±0.11 ^{NS}	2.42±0.07	2.42±0.07	1.99±0.04	2.24±0.02	2.37±0.04
	b*	5.95±0.88 ^{ac}	5.47±0.50 ^a	6.10±0.92 ^{ac}	7.91±0.88 ^{bd}	8.26±0.14 ^{bd}	7.22±0.18 ^{cd}
Crust	L*	64.28±2.51 ^a	62.85±1.65 ^a	63.53±1.30 ^a	70.93±1.90 ^b	77.01±1.51 ^c	78.85±0.76 ^c
	a*	14.68±1.17 ^a	16.08±0.63 ^a	14.75±0.75 ^a	7.28±1.89 ^b	3.36±0.24 ^b	3.03±0.10 ^b
	b*	29.44±0.42 ^a	30.04±0.51 ^a	28.88±0.71 ^a	23.68±2.66 ^b	13.41±1.25 ^c	11.19±0.83 ^c
	BI	75.89±4.81 ^a	81.22±2.05 ^a	75.46±3.12 ^a	47.23±8.61 ^b	21.69±0.84 ^d	17.59±1.10 ^d

The analysis of the color of white pan bread reveals the following: RC shows 0% B2, R1 reveals 20% B2, R2 presents 40% B2, R3 exhibits 60% B2, R4 indicates 80% B2, and R5 displays 100% B2.

^{a-d}Distinct letters placed above each column type signify a statistical difference at $\alpha=0.05$.

^{NS}Denotes non-significant results.

Table 4. TPA of Baromi2 White Pan Bread

	RC	R1	R2
Cohesiveness, 24 h (-)	0.69±0.00 ^{NS}	0.67±0.00	0.54±0.00
Springiness, 24 h (mm)	9.66±0.02 ^{NS}	9.64±0.01	9.61±0.02
Chewiness, 24 h (mJ)	24.66±0.08 ^a	24.79±0.07 ^a	34.15±0.27 ^b
Hardness, 24 h (N)	37.00±2.22 ^a	38.25±1.10 ^a	65.87±5.38 ^b
Hardness, 48 h (N)	45.47±1.10 ^a	52.67±5.83 ^b	75.15±3.91 ^c
Hardness, 96 h (N)	93.45±1.34 ^b	72.87±1.03 ^a	126.35±4.59 ^c

Texture of white pan bread: RC has 0% "Baromi2 (B2)"; R1 has 20% B2; R2 has 40%.

^{a-c}Above each column indicate statistical differences at $\alpha=0.05$.

^{NS}Nonsignificant. Letters.

ing hardness at 24 hr after baking, RC measured 37.00 N and R1 measured 38.25 N, while a sharp increase was noted for R2 at 65.87 N. After 48 hr, RC recorded a rise to 45.47 N (up 22.89% from the previous day), R1 reached 52.67 N (up 37.73%), and R2 saw a minimal increase to 75.15 N (1.08%). By 96 hr, RC jumped to 93.45 N (up 105.52% from the prior day), R1 rose to 72.87 N (up 38.37%), and R2 demonstrated a significant increase to 126.35 N (98.01%).

In general, aging occurred at a slower pace with increased B2 content. Even though R2 showed a reduced aging rate, its level was notably high. When R1 is compared to RC, the hardness increase rate saw a reduction of 22.01%. These findings imply that substituting 20% of B2 enhanced the aging rate and the overall aging extent. According to Nivell et al. (2020), the gluten network in bread influences the product's hardness, while the amylose network helps reduce it. This study concludes that substituting 20% of the wheat flour with B2 in baked goods reduced gluten content, which improved both the degree and rate of retrogradation, ultimately enhancing product quality.

Conclusion

In this study, we investigated the potential of using B2 as a wheat flour alternative to increase rice flour consumption in bread. We compared the physicochemical properties of products with added B2. The fermentation characteristics improved significantly when the B2 substitution rate was kept at optimal levels. Using B2 at a 20%

substitution in bread led to impressive fermentation expansion capabilities. The moisture content of the products decreased as the concentration of B2 increased. Color analysis showed that the BI value of the crust lowered with higher B2 levels, making the crust appear brighter. In texture evaluations, bread containing B2 showed reduced staling compared to the control group, with both hardness and staling rates declining at the 20% substitution. This study concludes that about 20% B2 was optimal for fermented bread products. However, thorough optimization is necessary to find the ideal B2 substitution rate and to mix various ingredients while upholding product quality. This research demonstrates the viability of using B2 as a wheat flour substitute, and it is expected that future studies involving different baking enzymes and gluten alternatives will further improve the quality of rice-enriched bread products.

ORCID

Ga Hyeon Kim	https://orcid.org/0009-0008-7311-8217
Sung Huo Kim	https://orcid.org/0009-0001-7644-7410
Yeon-Ji Jo	https://orcid.org/0000-0001-5226-3759
Sung Hoon Park	https://orcid.org/0000-0002-3032-5901

Conflict of interests

No potential conflict of interest relevant to this article was reported.

Acknowledgements

This work was carried out with the support of "Cooperative Research Program for Agriculture Science and Technology Development (RS-2022-RD010419)" Rural Development Administration, Republic of Korea.

Data availability

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authorship contribution statement

Conceptualization: Kim GH, Kim SH.
Data curation: Kim GH, Kim SH, Jo YJ.
Formal analysis: Kim GH, Kim SH.
Methodology: Kim GH, Kim SH, Jo YJ, Park SH.
Validation: Jo YJ, Park SH.

Investigation: Kim GH, Kim SH.

Writing - original draft: Kim GH, Kim SH.

Writing - review & editing: Kim GH, Kim SH, Jo YJ, Park SH.

Ethics approval

Not applicable.

References

- AACC. 2000. 10-10.03 Optimized Straight-Dough Bread-Making Method. AACC Off. Method.
- Aljada B, Zohni A, El-Matary W. 2021. The gluten-free diet for celiac disease and beyond. *Nutrients*, 13: 265.
- Baek JE, Koh BK. 2014. Bread and noodle making potential of commercial rice flour. *J. Living Sci. Res.* 40: 25-34.
- Caio G, Volta U, Sapone A, Leffler DA, De Giorgio R, Catassi C, Fasano A. 2019. Celiac disease: a comprehensive current review. *BMC Med.* 17: 142.
- Cannas M, Pulina S, Conte P, Del Caro A, Urgeghe PP, Piga A, Fadda C. 2020. Effect of the substitution of rice flour with quinoa flour on the chemical-physical, nutritional, volatile and sensory parameters of gluten-free ladyfinger biscuits. *Foods*, 9: 808.
- Choi SY, Lee YT. 2007. Properties of rice flour milled from spray-washed rice during storage. *J. Korean Soc. Food Sci. Nutr.* 36: 915-920.
- Chu JH, Choi JH, Go ES, Choi HY. 2023. Quality characteristics of the muffins prepared with different types of rice flour. *Korean J. Food Preserv.* 30: 630-641.
- Fujii K, Usui M, Ohsuga A, Tsuji M. 2023. Effect of thermoresponsive xyloglucan on the bread-making properties and preservation of gluten-free rice-flour bread. *Foods*. 121: 2761.
- Ha SK, Kim BK, Hwang WH, Mo Y, Jeong JM, Lee DK, Kim WJ, Kim JJ, Jeong JU. 2022. Early maturing rice variety “Baromi2” with a floury endosperm and suitable for dry-milling of rice grain. *Korean Soc. Breed. Sci.* 54: 433-441.
- Honda Y, Inoue N, Kurita M, Okunishi T. 2021. Alpha-glutelin degradation and its hydrolysate by protease enhance the specific volume of gluten-free rice starch bread. *J. Cereal Sci.* 102: 103338.
- Hu G, Burton C, Yang C. 2010. Efficient measurement of the amylose content in cereal grains. *J. Cereal Sci.* 51: 35-40.
- Jun HI, Yoo SH, Song GS, Kim YS. 2019. Bread-making properties of rice bread with naked oat flours. *Korean J. Food Preserv.* 26: 68-73.
- Kim MH. 2017. Finding a solution to the rice surplus problem. *Agric. Res.* 61: 140-155.
- Kim WM, Yoon KH, Lee GH. 2019. Physicochemical properties of gluten-free rice pan bread by adding processed rice flour. *J. Korean Soc. Food Sci. Nutr.* 48: 1143-1152.
- Kraithong S, Lee S, Rawdkuen S. 2018. Physicochemical and functional properties of Thai organic rice flour. *J. Cereal Sci.* 79: 259-266.
- Kupiec M, Zbikowska A, Marciniak-Lukasiak K, Zbikowska K, Kowalska M, Kowalska H, Rutkowska J. 2021. Study on the introduction of solid fat with a high content of unsaturated fatty acids to gluten-free muffins as a basis for designing food with higher health value. *Int. J. Mol. Sci.* 22: 9220.
- Mariscal-Moreno RM, Chuck-Hernández C, Figueroa-Cárdenas JD, Serna-Saldivar SO. 2021. Physicochemical and nutritional evaluation of bread incorporated with ayocote bean (*Phaseolus coccineus*) and black bean (*Phaseolus vulgaris*). *Processes*. 9: 1782.
- Monteiro JS, Farage P, Zandonadi RP, Botelho RBA, Oliveira L, Raposo A, Shakeel F, Alshehri S, Mahdi WA, Araújo WMC. 2021. A systematic review of gluten-free bread formulations using specific volume as a quality indicator. *Foods*. 10: 614.
- Nivelle MA, Beghin AS, Vrinten P, Nakamura T, Delcour JA. 2020. Amylose and amylopectin functionality during the storage of bread prepared from flour of wheat containing unique starches. *Food Chem.* 320: 126609.
- Park HY, Lee JH, Lee SJ, Kim Y, Kwak JE 2023. Analysis of the general ingredients and micronutrients in rice flour, including the new variety ‘Baromi2’, and wheat flour. *Korean J. Food Sci. Technol.* 55: 271-277.
- Qin W, Lin Z, Wang A, Xiao T, He Y, Chen Z, Wang L, Liu L, Wang F, Tong LT. 2021. Influence of damaged starch on the properties of rice flour and the quality attributes of gluten-free rice bread. *J. Cereal Sci.* 101: 103296.
- Roman L, Reguilon MP, Gomez M, Martinez MM. 2020. Intermediate-length amylose increases the crumb hardness of the rice flour gluten-free breads. *Food Hydrocolloids*. 100: 105451.
- Shewry P. 2019. What is gluten—why is it special? *Front. Nutr.* 6: 101.
- Šmídová Z, Rysová J. 2022. Gluten-free bread and bakery products technology. *Foods*, 11: 480.
- Sun H, Ju Q, Ma J, Chen J, Li Y, Yuan Y, Hu Y, Fujita K, Luan G. 2019. The effects of extruded corn flour on the rheological properties of the wheat-based composite dough and the bread quality. *Food Sci. Nutr.* 7: 2977-2985.

Woomer J, Singh M, Vijayakumar P, Adedeji A. 2019. Physical properties and organoleptic evaluation of gluten-free bread from proso millet. *Br. Food J.* 122: 547-560.

Zhao Q, Guo H, Hou D, Laraib Y, Xue Y, Shen Q. 2021. Influence of temperature on the storage characteristics of different rice varieties. *Cereal Chem.* 98: 935-945.